

Subfilter Scale Fluxes in the Marine Surface Layer

Peter P. Sullivan

National Center for Atmospheric Research
Boulder, CO 80307-3000

Phone:(303) 497-8953 fax:(303) 497-8171 email: pps@ncar.ucar.edu

James B. Edson

Department of Marine Sciences
University of Connecticut
Groton, CT 06340

Phone:(860) 405-9165 fax:(860) 405-9153 email: james.edson@uconn.edu

John C. Wyngaard

Department of Meteorology, Pennsylvania State University
627A Walker Building
University Park, PA 16802

Phone:(814) 863-7714 fax:(814) 865-3663 email: wyngaard@ems.psu.edu

Award #: NN0001406IP20018

<http://www.atd.ucar.edu/rtf/projects/OHATS04>

<http://www.whoi.edu/science/AOPE/dept/OHATS/intro.html>

LONG-TERM GOALS

A goal of marine surface layer research is to identify and quantify coupling mechanisms that connect the atmospheric boundary layer and surface waves. Large-eddy simulation (LES) plays a role in this research and has provided insight into the interactions between imposed waves and turbulence (Sullivan *et al.*, 2004; Sullivan *et al.*, 2006). However, the fidelity of subfilter-scale (SFS) parameterizations used in LES for flows over complex geometry, *e.g.*, a moving surface gravity wave field, is untested. Recent field campaigns such as the Horizontal Array Turbulence Study (HATS) conducted over land have provided new impetus to improve parameterizations in LES codes (*e.g.*, see Sullivan *et al.*, 2003; Chen & Tong, 2006; Hatlee & Wyngaard, 2006). A natural progression in these investigations is then to study increasingly complex flows. In the present work we present results from a new field campaign, the Ocean Horizontal Array Turbulence Study (OHATS), specifically directed at the measurement of SFS variables in the marine surface layer in the presence of surface waves. These observations can be used to examine and improve the SFS parameterization in LES codes, and more broadly, the impacts of water waves on surface layer turbulence under a variety of atmospheric stability conditions and wave states. Ultimately, the dataset and derived improvements to LES can benefit surface-layer parameterizations in mesoscale and numerical weather prediction models. Partners in OHATS are Applied Ocean Physics & Engineering at Woods Hole Oceanographic Institute (also U. Connecticut), the Earth Observing Laboratory at the National Center for Atmospheric Research, and the Department of Meteorology at Pennsylvania State University. The OHATS field campaign was conducted in 2004 off the coast of Martha's Vineyard using a low-profile Air-Sea Interaction Tower (ASIT). The ASIT is an integral component of the Martha's Vineyard Coastal Observatory operated by Woods Hole Oceanographic Institute. See <http://www.atd.ucar.edu/rtf/projects/OHATS04> for additional information.

| Report Documentation Page | | | | Form Approved OMB No. 0704-0188 | |
|--|------------------------------------|-------------------------------------|---|---|---------------------------------|
| Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| 1. REPORT DATE 30 SEP 2006 | | 2. REPORT TYPE | | 3. DATES COVERED 00-00-2006 to 00-00-2006 | |
| 4. TITLE AND SUBTITLE Subfilter Scale Fluxes in the Marine Surface Layer | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Center for Atmospheric Research,Box 3000,Boulder,CO,80307-3000 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT Same as Report (SAR) | 18. NUMBER OF PAGES 9 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

OBJECTIVES

The immediate goal of our research is to carry out an extensive analysis of the large database collected during OHATS. Bulk and SFS variables (*e.g.*, momentum and scalar fluxes) will be computed as function of atmospheric stratification and wave age. These findings can then readily be compared with previous results obtained over land.

APPROACH

Large-eddy simulation faces a closure problem similar to that in traditional Reynolds average modeling (RANS) (*e.g.*, see Pope, 2000). In LES the unknowns are the SFS fluxes for momentum, defined as

$$\tau_{ij} = \overline{u_i u_j} - \overline{u_i} \overline{u_j} , \quad (1)$$

and the SFS flux for an arbitrary scalar c defined as

$$f_i = \overline{u_i c} - \overline{u_i} \overline{c} . \quad (2)$$

In the above the overbar $\overline{(\)}$ denotes *spatial* filtering.

The scientific approach adopted here is to obtain field measurements of the total fields (u_i, c) from which we can construct SFS momentum and scalar fluxes in (1) and (2) over a wide range of stratification and wave states. The data needs to be gathered at a sufficient number of spatial locations to permit spatial filtering of the total fields.

OHATS utilizes the horizontal array technique first proposed by Tong *et al.* (1998, 1999) which has since been utilized in several investigations of SFS motions in atmospheric boundary layers over rough land surfaces (Porté-Agel *et al.*, 2001; Horst *et al.*, 2004; Sullivan *et al.*, 2003). Two arrays (or lines) of sonic anemometers are positioned crosswind to the primary wind direction with the horizontal spacing between individual sonic anemometers selected to achieve a specific spatial filter width. The two sonic lines are located at different heights above the surface to allow for measurement of vertical gradients. Under the assumption of Taylor's hypothesis in the alongwind direction, we are in essence able to measure a 2-D ($x - y$) plane of winds and temperature fluctuations at two levels in the atmospheric surface layer. The use of a dense array of sonic anemometers allows both single and double spatial filtering. In addition to the surface layer winds in OHATS, wave height information and wave propagation direction were obtained using three downward pointing laser altimeters mounted on the ASIT diving board.

WORK COMPLETED

During fiscal year 2005 we completed the deployment for OHATS and successfully carried out the field campaign. The intensive observation period for OHATS extended from August to October, approximately 85 days. A description of the data collection procedures, the available data, and software tools used to analyze the database are described in our FY05 progress report. In FY06 we began a detailed analysis of the OHATS data. This includes computation of bulk surface layer variables, SFS momentum and scalar fluxes, and transfer rates between resolved and SFS fields. Also, wave height spectra were computed using data collected from the laser altimeters. The availability of the wave height information provides an opportunity to correlate motions in the air with the underlying sea surface. The wave-induced motion algorithm used in OHATS was provided by Dr. Tihomir Hristov (Johns Hopkins University) and is described in Hristov *et al.* (1998). An overview of the data analysis procedures and software is given in Sullivan *et al.* (2006).

RESULTS

The OHATS database provides an opportunity to perform a variety of analysis on SFS variables and the underlying wave fields. Recently, we analyzed the SFS momentum and scalar fluxes with a goal of evaluating different SFS parameterizations. Here we concentrate on the the SFS scalar flux and examine the similarities and differences for flow over stationary roughness and moving water waves.

Wyngaard (2004) proposed a new class of subfilter-scale models that are intended to be applicable across a range of scales spanning the gap $l \sim \Delta_f$ between the “LES limit” $l/\Delta_f \gg 1$ and the “mesoscale limit” $l/\Delta_f \ll 1$ with l the scale of the dominant turbulence and Δ_f the filter width. The mathematical steps outlining the methodology are fully described by Wyngaard (2004). The basis of these simplified rate-equation models is rational truncation of the full transport equations for SFS fluxes based on scale analysis. These SFS prescriptions offer improvements to simple eddy viscosity closures, but have not yet been fully implemented and evaluated in simulation codes. Using a subset of the HATS database (Sullivan *et al.*, 2003), Hatlee & Wyngaard (2006) tested variants of these SFS closures for scalar flux. One of the important results from their investigation is the proper prediction of all components of SFS scalar flux. For a conserved scalar c the components of the SFS scalar flux f_i are modeled using a rate equation of the form

$$\frac{\partial f_i}{\partial t} = -f_j \frac{\partial \bar{u}_i}{\partial x_j} - \tau_{ij} \frac{\partial \bar{c}}{\partial x_j} - \frac{f_i}{T} . \quad (3)$$

This truncated model includes only time change, tilting and production terms and models pressure destruction via a sink of the form $-f_i/T$ with the time scale of the SFS turbulence

$$T = C \Delta_f / E^{1/2} . \quad (4)$$

In the above expressions, \bar{u}_i is the resolved (filtered) velocity, τ_{ij} is the SFS momentum flux, $E = \tau_{ii}/2$ is the SFS kinetic energy, and the modeling constant $C \approx 0.3$ is chosen to match the HATS data.

Algebraic stress models with some structural similarity to (3) have been proposed for stratified flows (Findikakis & Street, 1979) and implemented in RANS codes (*e.g.*, Hanjalic, 2001), but only recently have these types of models been validated against geophysical data in the context of LES. Here we test the applicability of (3) and (4) for scalar transport in turbulent flow over water waves.

The OHATS database possesses all the ingredients necessary to evaluate the SFS closure given by (3) using virtual temperature θ as the conserved scalar c . Our algorithm advances (3) in time with a second-order Adams-Bashforth method. The observed values of SFS momentum fluxes and energy, resolved velocity and scalar gradients, which appear on the right-hand-side, are spline interpolated to increase their temporal resolution and thereby eliminate errors due to finite differencing. We performed an independent test of our algorithm using all the acceptable cases in the HATS database. Figure 1 shows the observed and modeled components of average scalar flux $\langle f_1 \rangle, \langle f_3 \rangle$; these results are in agreement with Hatlee & Wyngaard (2006) and verify the correctness of our algorithm. The findings illustrate that (3) is a good predictor for SFS scalar flux over a range of stratifications and filter widths over land. An important attribute of (3) is the prediction of a finite average value of horizontal scalar flux $\langle f_1 \rangle$ even in a horizontally homogeneous flow $\langle \partial \bar{c} / \partial x, \partial \bar{c} / \partial y \rangle = 0$. An eddy-viscosity scalar-gradient closure leads to $\langle f_1 \rangle = 0$. We mention that a dynamic evaluation of the eddy viscosity (*e.g.*, Germano *et al.*, 1991) cannot overcome this deficiency since production from “scalar tilting” $f_j \partial \bar{u}_i / \partial x_j$ in (3) is not present in eddy viscosity models (see also Wyngaard *et al.*, 1971).

Results from our analysis of the OHATS data are depicted in figure 2. These findings show that the prediction of horizontal scalar flux over waves is slightly below but closely follows the trend from HATS. However, (3) noticeably over predicts the SFS vertical scalar flux f_3 especially in cases with larger waves and scalar surface flux. Cases with significant wave heights greater than 1 m and at the same time surface convection exhibit the greatest departure from the HATS observations and predictions given by (3). The trends in figure 2 seem clear and hence the surprising result for f_3 merits further interrogation.

Inspection of (3) suggests the approximate balance $\tau_{ij}\partial\bar{c}/\partial x_j \approx -f_i/T$, *i.e.*, the production of SFS flux is balanced by pressure destruction. Then a simple (*ad hoc*) modification is to introduce separate pressure destruction modeling constants (C_H, C_V) in the horizontal and vertical directions, respectively. The physical motivation for this model is the speculation that the wave field modifies the production of SFS scalar flux and hence the pressure destruction needs to respond accordingly. A test of this modified scalar flux model is shown in figure 3 using best constants $(C_H, C_V) = (0.3, 0.06)$. The model predictions closely follow the observations; note the predictions with this modified model improves both the horizontal and vertical scalar flux estimates. An important ingredient of any SFS model is to transfer squared scalar variance \bar{c}^2 between resolved and SFS fields, *i.e.*, establish a proper cascade to smaller scales. The variance transfer rate for scalars is (Wyngaard, 2004)

$$\mathcal{P} = 2 f_i \frac{\partial \bar{c}}{\partial x_i} . \quad (5)$$

In the LES regime, \mathcal{P} is a stochastic variable and not constrained to be positive definite, *i.e.*, the instantaneous transfer can be from the SFS field to the resolved field (backscatter) or from the resolved field to the SFS field (forwardscatter). However, the ensemble average $\langle \mathcal{P} \rangle$ is forwardscatter. In figure 4 we compare estimates of mean scalar variance transfer from our model to the OHATS observations. The close agreement indicates that the modified rate equation (with 2 pressure destruction constants) performs well using the scalar transfer rate as a metric.

These results for SFS scalar flux suggest that the wave field plays a role in setting the SFS fluxes that is unique compared to flow over stationary roughness. A next step is to closely examine the phase relationship between the wave induced pressure and scalar fields. In order to explore this speculation we need to identify the wave correlated motions in the velocity, pressure, and scalar fields, as described earlier. In addition, we can use the direct numerical simulation (DNS) results for stratified flow over waves described by Sullivan & McWilliams (2002) to gain insight into the present results.

IMPACT/APPLICATIONS

The OHATS field enhances our current understanding of air-sea interaction processes, provides insight into the couplings between surface waves and turbulence, and sheds new light on SFS modeling for LES. The research also compliments the CBLAST program sponsored by the Office of Naval Research. In addition the OHATS dataset can be used to further our understanding of LES of swell driven marine boundary layers (Sullivan *et al.*, 2004; Sullivan *et al.*, 2006; Edson *et al.*, 2006).

TRANSITIONS & RELATED PROJECTS

We are currently engaged in using large-eddy simulation to model air-sea interaction processes in both the atmospheric and oceanic boundary layers. In particular, we are using LES to assist in

the interpretation of field measurements taken during the recent low-wind (Edson *et al.*, 2006) and hurricane Coupled Boundary Layer Air-Sea interaction Transfer (CBLAST) field campaigns. OHATS provides an additional data source for understanding air-sea interaction processes. A web site describing CBLAST and the Martha's Vineyard Coastal Observatory can be found at:

<http://www.whoi.edu/science/AOPE/dept/CBLASTmain.html>

<http://mvcodata.whoi.edu/cgi-bin/mvco/mvco.cgi>

REFERENCES

- Chen, Q. & C. Tong, 2006: Investigation of the subgrid-scale stress and its production rate in a convective atmospheric boundary layer using measurement data. *J. Fluid Mechanics*, **547**, 65-104.
- Edson, J., T. Crawford, J. Crescenti, T. Farrar, J. French, N. Frew, G. Gerbi, C. Helmis, T. Hristov, D. Khelif, A. Jessup, H. Jonsson, M. Li, L. Mahrt, W. McGillis, A. Plueddmann, L. Shen, E. Skillingstad, T. Stanton, P. Sullivan, J. Sun, J. Trowbridge, D. Vickers, S. Wang, Q. Wang, R. Weller, J. Wilkin, D. Yue, & C. Zappa, 2006: The coupled boundary layers and air-sea transfer experiment in low winds (CBLAST-Low). *Bulletin of the American Meteorological Society*, accepted.
- Findikakis, A.N. & R.L. Street, 1979: An algebraic model for subgrid-scale turbulence in stratified flows. *J. Atmospheric Sciences*, **36**, 1934-1949.
- Germano, M., U. Piomelli, P. Moin, & W.H. Cabot, 1991: A dynamic subgrid-scale eddy viscosity. *Physics of Fluids*, **3**, 1760-1765.
- Hanjalic, K. & S. Kenjeres, 2001: VLES of flows driven by thermal buoyancy and magnetic field. In *Modern Simulation Strategies for Turbulent Flow*, ed. B.J. Geurts, publisher R.T. Edwards, pp. 223-246.
- Hatlee, S.C. & J.C. Wyngaard, 2006: Subfilter-scale modeling in the Terra Incognita. *J. Atmos. Sci.*, **submitted**.
- Horst, T. W., J. Kleissl, D. H. Lenschow, C. Meneveau, C-H. Moeng, M. B. Parlange, P. P. Sullivan & J. C. Weil, 2004: Field observations to obtain spatially-filtered turbulence fields from transverse arrays of sonic anemometers in the atmospheric surface layer, *J. Atmos. Sci.*, **61**, 1566-1581.
- Hristov, T., C. Friehe & S. Miller, 1998: Wave-coherent fields in air flow over ocean waves: Identification of cooperative behavior buried in turbulence, *Physical Review Letters*, **81**, 5245-5248.
- Pope, S. B., 2000: *Turbulent Flows*, Cambridge University Press, pp. 771.
- Porté-Agel, F., M.B. Parlange, C. Meneveau, & W. E. Eichinger, 2001: A priori field study of the subgrid-scale heat fluxes and dissipation in the atmospheric surface layer, *J. Atmos. Sci.*, **58**, 2673-2698.
- Sullivan, P.P, T.W. Horst, D.H. Lenschow, C-H. Moeng, & J.C. Weil, 2003: Structure of subfilter-scale fluxes in the atmospheric surface layer with application to large-eddy simulation modeling. *J. Fluid Mech.*, **482**, 101-139.
- Sullivan, P.P., J.B. Edson, J.C. McWilliams, C-H. Moeng, 2004: Large-Eddy Simulations and Observations of Wave Driven Boundary Layers. *16th Symposium on Boundary Layer and Turbulence*, Portland, ME.

- Sullivan, P. P., J. B. Edson, T. Hristov, & J. C. McWilliams, 2006: Momentum flux structures and statistics in low-wind marine surface layers: Observations and large-eddy simulations. *27th Conference on Hurricanes and Tropical Meteorology*, Monterey, CA.
- Sullivan, P.P. & J.C. McWilliams, 2002: Turbulent flow over water waves in the presence of stratification, *Physics of Fluids*, **14**, 1182-1195.
- Tong, C., J.C. Wyngaard, S. Khanna, & J.G. Brasseur, 1998: Resolvable- and subgrid-scale measurement in the atmospheric surface layer: Technique and issues. *J. of Atmos. Sci.*, **55**, 3114-3126.
- Tong, C., J.C. Wyngaard, & J.G. Brasseur, 1999: Experimental study of the subgrid-scale stresses in the atmospheric surface layer.
- Wyngaard, J.C., O. R. Côte, & Y. Izumi, 1971: Local free convection, similarity, and the budgets of shear stress and heat flux. *J. Atmos. Sci.*, **28**, 1171-1182.
- Wyngaard, J.C., 2004: Toward numerical modeling in the "Terra Incognita". *J. Atmos. Sci.*, **61**, 1816-1826.

PUBLICATIONS

- Beare, R. J., M. K. Macvean, A. A. M. Holtslag, J. Cuxart, I. Esau, J-C. Golaz, M. A. Jimenez, M. Khairoutdinov, B. Kosovic, D. Lewellen, T. S. Lund, J. K. Lundquist, A. McCabe, A. F. Moene, Y. Noh, S. Raasch & P. P. Sullivan, 2006: An intercomparison of large-eddy simulations of the stable boundary layer. *Boundary-Layer Meteorology*, **118**, 242-272.
- Edson, J., T. Crawford, J. Crescenti, T. Farrar, J. French, N. Frew, G. Gerbi, C. Helmis, T. Hristov, D. Khelif, A. Jessup, H. Jonsson, M. Li, L. Mahrt, W. McGillis, A. Plueddmann, L. Shen, E. Skillingstad, T. Stanton, P. Sullivan, J. Sun, J. Trowbridge, D. Vickers, S. Wang, Q. Wang, R. Weller, J. Wilkin, D. Yue, & C. Zappa, 2006: The coupled boundary layers and air-sea transfer experiment in low winds (CBLAST-Low). *Bulletin of the American Meteorological Society*, accepted.
- Edson, J., C. Fairall, & P. Sullivan, 2006: Evaluation and continued improvements to the TOGA COARE 3.0 algorithm using CBLAST data. *27th Conference on Hurricanes and Tropical Meteorology*, Monterey, CA.
- Moeng, C.-H. J. Dudhia, J. Klemp & P. Sullivan, 2006: Examining two-way grid nesting for large-eddy simulation of the PBL using the WRF model. *Monthly Weather Review*, submitted.
- Patton, E., P. Sullivan, K. Ayotte, & J. Finnigan, 2006: Turbulent flow over isolated ridges: Influence of vegetation. *17th Conference on Boundary Layer and Turbulence*, San Diego, CA.
- Patton, E. G., P. P. Sullivan, & K. W. Ayotte, 2006: Flow and transport above and within forests in complex topography. Preprints *1st Integrated Land Ecosystem - Atmospheric Processes Study* (iLeaps) Scientific Conference, Jan. 23-26, Boulder, CO. Report series In Aerosol Science, No. 79, Editors: A Reissell and A. Aarflot, pp 283-284.
- Sullivan, P.P., J.C. McWilliams, W.K. Melville, 2006: Surface gravity wave effects in the oceanic boundary layer: Large-eddy simulation with vortex force and stochastic breakers. *Journal of Fluid Mechanics*, submitted.
- Sullivan, P. P., J. B. Edson, J. C. Wyngaard, & M. Kelly, 2006: Subfilter scale fluxes in the marine surface layer: Results from the Ocean Horizontal Array Turbulence Study (OHATS). *17th Conference on Boundary Layer and Turbulence*, San Diego, CA.

Sullivan, P. P., J. B. Edson, T. Hristov, & J. C. McWilliams, 2006: Momentum flux structures and statistics in low-wind marine surface layers: Observations and large-eddy simulations. *27th Conference on Hurricanes and Tropical Meteorology*, Monterey, CA.

Weil, J. C., E. G. Patton, & P. P. Sullivan, 2006: Lagrangian particle modeling of dispersion in the stable boundary layer. *17th Conference on Boundary Layer and Turbulence*, San Diego, CA.




Figure 1: Comparison of modeled and observed SFS scalar flux in the atmospheric surface layer over a stationary rough surface from the field campaign HATS (Sullivan *et al.* 2003). $\langle f_1 \rangle$ and $\langle f_3 \rangle$ are indicated by black and red dots respectively. The inset photograph shows the HATS field deployment of an array of sonic anemometers above a fallow field in California.




Figure 2: Comparison of modeled and observed SFS scalar flux in the marine surface layer over moving waves. Horizontal scalar flux $\langle f_1 \rangle$ is indicated by open blue circles. Vertical scalar flux $\langle f_3 \rangle$ is indicated by open green and orange circles; cases with significant wave height greater than 1 m are shown in orange. The results from HATS are the black and red dots. Note the difference in vertical and horizontal scales compared to figure 1. The inset photograph taken from the ASIT shows sonic anemometers before final positioning in OHATS.




Figure 3: Comparison of modeled and observed SFS scalar flux in the marine surface layer over moving waves as in figure 2 but using a modified rate equation with separate constants in the pressure destruction model for horizontal and vertical scalar flux, $(C_H, C_V) = (0.3, 0.06)$, respectively.




Figure 4: Comparison of modeled and observed mean scalar transfer $\langle f_i \frac{\partial \bar{\theta}}{\partial x_i} \rangle$ in the marine surface layer over moving waves. Orange dots indicate results for cases with significant wave heights greater than 1m and convective conditions. Blue dots are all other cases.